

## **Colonoscopy Trilyte (Colyte, Golyte, Nulyte) EXTENDED Split Dose**

**\* IF TAKING PHENTERMINE (Adipex-P) – MUST STOP 7 DAYS BEFORE YOUR PROCEDURE\***

### **FIVE DAYS BEFORE EXAM:**

- 1) Discontinue any Vitamin E, herbal medications, Fish Oil, Fiber or Iron Supplements.
- 2) Do not eat any corn, nuts, seeds or foods containing seeds (Example: Watermelon, Squash, Cucumbers, Tomatoes, Rye Bread)

### **TWO DAYS BEFORE EXAM:**

- 1) At 5pm **No Solid Foods**–Have full liquid dinner includes: Shakes, Smoothies & Yogurt without chunks of fruit or Seeds. Or Clear liquids are; Gatorade or PowerAde, Water, Clear fruit juices, Clear sodas, Bouillon or Chicken broth, plain Jello, Popsicles. You may have plain black coffee, tea.

### **ONE DAY BEFORE EXAM:**

- 1) If on any Blood Pressure or heart medications, please take your normal dose today.
- 2) *Diabetic Patients should take ½ of your usual dose today. \*Note –Patients with an insulin pump contact your PCP/Endocrinologist on how to adjust your pump settings when fasting.*
- 3) Discontinue Alcohol one day prior to exam.
- 4) Drink only clear liquids for breakfast, lunch and dinner the entire day. **Clear liquids allowed; You may have plain black coffee, tea, water, clear fruit juices, clear sodas, bouillon or plain chicken broth, plain jell-O, or popsicle., Only Orange or Yellow colors (No Red, Purple, Blue or Green Color Dyes).**

#### **NO SOLID FOODS ALL DAY, NO MILK OR MILK PRODUCTS ALLOWED**

**HYDRATION** is Very Important to the procedure – Drinking 8oz of clear liquids every hour throughout the day before you start taking the prep should be your goal in order to be properly hydrated.

- 5) Prepare **Trilyte or (Colyte, Golyte, Nulyte)** Add tap water to FILL line. Replace cap tightly and mix or shake well until ingredients have dissolved. No other ingredients should be added to the solution. Refrigerate until ready to drink. Chilling improves the palatability (store no longer than 48 hours)
- 6) **(First Dose)** At 6:00pm Begin drinking ½ the solution (2 Liters), Drink an 8-oz. glass every 10 minutes. It is best to drink the whole glass rapidly, rather than sipping small amounts. Bowel movements should begin approximately 1 hour after the first glass and will continue periodically for several hours.
- 7) You are strongly encouraged to drink as much clear liquids as you want until you go to sleep.

### **DAY OF EXAM:**

1. **No Solid Foods, No Milk or Milk Products**
2. **(Second Dose) Exactly 7 hours before your procedure:** Start Time: \_\_\_\_\_ Finish Time: \_\_\_\_\_  
Start drinking the remaining solution (2 Liters) an 8-oz glass every 10 minutes until gone.  
**Then nothing to eat, drink, or chew (including water) until after your procedure is done. The prep must be finished 6 hours prior to the procedure. Failure to do so may require extra wait time after you arrive or could possibly result in cancellation & rescheduling of the procedure.**
3. **TRANSPORTATION REQUIRED**- you will not be able to drive after the procedure.
4. If you are a smoker, for anesthesia requirements do not smoke the day of the procedure.
5. If on any Blood Pressure or Heart Medications, please take your normal dosage in the early A.M.- 1<sup>st</sup> thing upon waking up- with a sip of water. Nothing else to eat, drink or chew until after your procedure.
6. Diabetics – Do NOT take your insulin or diabetic pills the day of the procedure.

### **YOUR PROCEDURE IS SCHEDULED ON:**

DATE: \_\_\_\_\_ TIME: \_\_\_\_\_ BE HERE AT: \_\_\_\_\_  
PLACE: Barkley Surgicenter 63 Barkley Circle Suite # 104 Ft. Myers, Florida 33907

### **YOUR POST PROCEDURE OFFICE VISIT IS SCHEDULED ON:**

DATE: \_\_\_\_\_ TIME: \_\_\_\_\_ WITH: \_\_\_\_\_  
 Ft. Myers 4790 Barkley Cir Bldg. A  Cape Coral 1303 SE 8<sup>th</sup> Terr.  Bonita 3501 Health Center Blvd. Ste 2130  
**If you have any questions regarding the procedure, call our office at (239) 275-8882, (239) 458-0822 or Visit our website at [www.giaswfl.com](http://www.giaswfl.com)**

**COLONOSCOPY**  
**Some Things You Should Know.....**

**WHAT IS A COLONOSCOPY?**

Colonoscopy is an examination of the large intestine using a long flexible tube called a colonoscope. This test may be done for a variety of reasons. Commonly it is done to investigate the finding of blood in the stool, diarrhea, abdominal pain, a change in bowel habits, or an abnormality noted on a colon x-ray or barium enema.

**WHAT HAPPENS DURING THE PROCEDURE?**

An intravenous (IV) will be started in your arm. Enough medication will be given to make you drowsy and relaxed.

The examination is performed with you lying on your left side. The doctor will examine your rectum first with a finger and then the colonoscope will be gently inserted. The colonoscope will be carefully passed through the colon and then slowly withdrawn.

If any abnormalities are found, small pieces of tissue may be taken for examination. Small growths (polyps) may also be found in the colon and can safely be removed. Small bleeding sites are often cauterized (burned) to stop bleeding. This is a very safe procedure and in many cases eliminates the need for surgery.

Your stay at the facility will be approximately three hours long, depending on how long it takes to pass the scope around the colon. When medication is given, the procedure is usually not unpleasant and often the patient does not remember having the procedure done.

**AFTER THE PROCEDURE:**

You will be observed for about thirty minutes before going home. You will not be allowed to drive after the procedure. You will be drowsy several hours afterward. Following the colonoscopy, the doctor will talk with you and your family and give them a report of the test findings. A follow-up visit should have been pre-scheduled so the test results can be explained to you when you are fully awake at a later date. This follow-up visit is separate from the procedure and if your insurance requires a co-pay it will be due on the day of the follow-up visit.

**RISKS:**

This is a very safe examination, but there is a small risk of complications, which occur in less than 1% of patients. Reactions to medications may occur. These might be allergic reactions or difficulty breathing. Bleeding is a possibility. Perforation of the colon may occur. A perforation is a serious problem and in some cases would require surgery to close the perforation of the colon.

If you experience severe pain, bleeding, or any problems after the test, please contact the doctor at (239) 275-8882 immediately.

Gastroenterology Associates of S.W. Florida, PA complies with applicable Federal civil rights laws and does not discriminate on the basis of race, color, national origin, age, disability, or sex.